


# 膀胱全摘除術後の成績

1996 - 2011	All	non-Neobladder	Neobladder	<i>P value</i>
n	299	130	169	
Age	67.2±9.4	70.3±9.8	66.5±11.2	<i>&lt;0.001</i>
Gender (M/F)	239 / 60	110 / 20	129 / 40	<i>n.s.</i>
CVD	42	22	20	<i>n.s.</i>
HT	119	55	64	<i>n.s.</i>
DM	35	16	19	<i>n.s.</i>
Stage	2.1±1.3	2.4±1.2	2.1±1.2	<i>&lt;0.001</i>
Complications	66	9	57	<i>&lt;0.001</i>
Recurrence	91	53	38	<i>&lt;0.001</i>
eGFR				
before	66.7±18.8	59.9±19.1	72.0±16.9	
1M	63.2±20.4	59.0±20.7	66.5±19.7	
1Y	62.7±19.2	57.6±22.3	65.7±16.6	
2Y	60.7±20.0	54.4±21.7	60.9±18.4	
3Y	61.1±23.6	53.8±28.2	65.2±19.6	
5Y	58.4±19.5	49.9±24.2	62.6±15.3	
Rates				
1M / Before	0.97±0.27	1.01±0.24	0.94±0.28	<i>0.037</i>
5Y / 1M	0.85±0.26	0.82±0.32	0.86±0.23	<i>n.s.</i>


# 膀胱癌にて膀胱全摘除術後の予後

Survival of BCa pts all n=299



- Stage 0, n=34      Stage 0; pT0
- Stage I, n=69      Stage I ; pT1
- Stage II, n=92      Stage 2; pT2
- Stage III, n=55      Stage 3; pT3~4a
- Stage IV, n=50      Stage 4; pT4b, N+, M+

Survival of Bladder Cancer pts, n=299



2群間に年齢差(約4歳)があるため、Neobladder群が長生きしやすい、と単純には言えない。